

The Charm Contribution to the Structure Function of the Proton

Ekaterini Tzamariudaki
MPI Munich



for the



and

collaborations


Outline of the Presentation

- Introduction
- D^* cross sections
- F_2^c
- Summary

abstracts: 853, 858, 983

Kinematics of Deep Inelastic Scattering

$27.5 \text{ GeV } e^\pm \rightarrow \sqrt{s} = 300 \text{ GeV} \leftarrow p 820 \text{ GeV}$



$$Q^2 = -q^2 = -(k - k')^2 \quad \text{4-momentum transfer}^2$$

$$x = Q^2 / 2P \cdot q \quad \text{fraction of } p \text{ momentum carried by the struck quark}$$


$$y = (p \cdot q) / (p \cdot k) \quad \text{relative energy transfer in the } p \text{ rest frame}$$

$$W^2 = (q + P)^2 \quad \text{mass}^2 \text{ of the hadronic final state}$$

$$Q^2 = sxy$$

$$\eta = -\ln(\tan \frac{\theta}{2})$$

Charm Production



- charm production dominated by the Boson Gluon Fusion mechanism
- sensitive to the gluon density

Hard scale allows validity of pQCD calculations

Various calculation schemes based on

- the DGLAP evolution

(Fixed Flavour Number Scheme (FFNS) massive and massless approach, Variable Flavour Number Scheme (VFNS))

- the CCFM evolution

(Ciafaloni, Catani, Fiorani and Marchesini)

Test of universality of the gluon distribution

Charm Tagging in DIS

- D^{*+} mesons



H1 and ZEUS




ZEUS

- semileptonic decays of charm into electrons

ZEUS


ZEUS Preliminary 1996-97



definition of “visible” range in terms of

$$p_t(D^{*+}, e) \quad \text{and} \quad \eta(D^{*+}, e)$$

Inclusive D^* Cross Sections



kinematic region

$1 < Q^2 < 100 \text{ GeV}^2$

$0.05 < y < 0.7$

$p_T D^* > 1.5 \text{ GeV}/c$

$|\eta_{D^*}| < 1.5$

H1

$$\sigma(e^+ p \rightarrow e^+ D^{*\pm} X) =$$

$$8.37 \pm 0.41(\text{stat.})^{+1.11}_{-0.82}(\text{syst.})^{+0.64}_{-0.39}(\text{theo.}) \text{ nb}$$

ZEUS


$1 < Q^2 < 600 \text{ GeV}^2, \quad 0.02 < y < 0.7$

$1.5 < p_T D^* < 15 \text{ GeV}/c, \quad |\eta_{D^*}| < 1.5$

$$\sigma(e^+ p \rightarrow e^+ D^{*\pm} X) = 8.31 \pm 0.31(\text{stat.})^{+0.3}_{-0.5}(\text{syst.}) \text{ nb}$$

Differential Cross Sections


ZEUS Data compared with HVQDIS



$$x_{D^*} = \left(\frac{2|\vec{p}_{D^*}|}{W} \right)$$

HVQDIS predictions for $m_c = 1.3 - 1.5 \text{ GeV}$, using the ZEUS 1994 PDF $p_t(c \rightarrow D^*)$, drag effect correction

Differential Cross Sections




HVQDIS and **CASCADE**

GRV98HO, $m_c 1.3 - 1.5 \text{ GeV}$
 $p_t(c \rightarrow D^*)$, $\epsilon 0.035 - 0.1$

gluon from F_2 fit, $m_c 1.3 - 1.5 \text{ GeV}$

Double Differential Cross Sections

H1 preliminary




Differential Cross Sections

$c \rightarrow e\nu X$

$1 < Q^2 < 1000 \text{ GeV}^2, \quad 0.03 < y < 0.7$

$1.2 < p_e < 5.0 \text{ GeV}, \quad 0.65 < \theta_e < 2.5 \text{ rad}$

ZEUS Preliminary 1996-97



Extraction of $F_2^c(x, Q^2)$

The $c\bar{c}$ production cross section described by $F_2^{c\bar{c}}(x, Q^2)$

$$\frac{d^2\sigma^{c\bar{c}}}{dxdQ^2} = \frac{2\pi\alpha^2}{xQ^4}(1 + (1 - y)^2) \cdot F_2^{c\bar{c}}(x, Q^2)$$


assumptions $F_L^c \sim 0$

extrapolate outside the “visible” region in p_T and η


$$F_2^{c \ meas}(x, Q^2) = \sigma_{vis}^{meas}(x, Q^2) \cdot \frac{F_2^{c \ theo}(x, Q^2)}{\sigma_{vis}^{theo}(x, Q^2)}$$

$F_2^c(x, Q^2)$

F_2^c in the NLO DGLAP scheme




$F_2^c(x, Q^2)$



- Strong Scaling violations

$$F_2^c(x, Q^2)/F_2(x, Q^2)$$


ZEUS Preliminary 1996-97




- Ratio $F_2^{c\bar{c}} / F_2$ is large $\sim 25 - 30\%$

$$F_2^c(x, Q^2)$$

F_2^c in the CCFM scheme



$F_2^c(x, Q^2)$



F_2^c extracted with HVQDIS
 F_2^c extracted with CASCADE

Conclusions

- D^* production cross sections
 - measured with improved precision
 - also double differentially
 - compared to NLO calculations

- F_2^{charm}
 - strong rise to lower x for increasing Q^2
 - large scaling violations
 - F_2^{charm}/F_2 reaches $\sim 30\%$ at low x and high Q^2

Charm Tagging in DIS

charm tagged in DIS using

- D^{*+} mesons



- semileptonic decays of charm into electrons ZEUS

definition of “visible” range in terms of

$$p_t(D^{*+}, e) \quad \text{and} \quad \eta(D^{*+}, e)$$

Luminosity

ZEUS: 37 pb^{-1} (34 pb^{-1}) for the D^{*+} (semileptonic) analysis
 H1: 18.6 pb^{-1}