

Salvatore Mele

INFN/Napoli CERN/EP



ZZ Cross-Section Measurements

On behalf of the four LEP Collaborations


- Signal Definition
- Physics Motivations
- Data Sample
- Analysis Procedure
- Results (also on $Z\gamma^*$)
- (Some) Interpretations

ICHEP2000

XXXth International Conference on High Energy Physics
Osaka, Japan, July 27th – August 2nd 2000



Signal Definition






- **Test of the Standard Model in the sector of Neutral Currents**
- **Study of possible Triple Vertices of Neutral Gauge Bosons (forbidden in the Standard Model) \Rightarrow Clara Matteuzzi's talk**
- **Sensitive to manifestations of new physics like Extra Spatial Dimensions (Low Scale Gravity)**
- **Also sensitive to excited electrons and SUSY, not yet studied experimentally**
- **Test Bench of the LEP detector capability to detect Higgs bosons**

Predictions, data sample and signatures

$\sim 0.5 \text{ fb}^{-1}/\text{experiment}$ $\sim 2.2 \text{ fb}^{-1}$ total



All ZZ results above 190 GeV are preliminary


Channel	Statistics	Signature	Background	Contamination
$q\bar{q}q'\bar{q}'$	High	4 jets	QCD $W^+W^- \rightarrow 4 \text{ jets}$	High
$q\bar{q}\nu\bar{\nu}$	High	2 jets + Missing energy	QCD $W^+W^- \rightarrow q\bar{q}l\nu$ $W\ell\nu$	Average
$q\bar{q}l^+l^-$	Average	2 jets + 2 leptons	W^+W^- 4f not ZZ	Low
$l^+l^-\nu\bar{\nu}$	Average	2 leptons + Missing energy	4f not ZZ W^+W^- l^+l^-	Average
$l_1^+l_1^-l_2^+l_2^-$	Low	4 leptons	4f not ZZ	Low



$$e^+e^- \rightarrow ZZ \rightarrow q\bar{q}e^+e^-$$

$M_{ee} = 93 \text{ GeV}$
 $M_{qq} = 89 \text{ GeV}$
 $M_{5C} = 91 \text{ GeV}$


L3

 $\sqrt{s} = 183 \text{ GeV}$ 



$$e^+e^- \rightarrow ZZ \rightarrow q\bar{q}l^+l^-$$

DELPHI Preliminary



- Invariant masses of leptons and hadrons
- Opening angles

Removing one of the mass constraints for $\mu^+\mu^-q\bar{q}$:

$$\sigma_{e^+e^- \rightarrow Z\gamma^* \rightarrow \mu^+\mu^-q\bar{q}}^{183 \text{ GeV} - 208 \text{ GeV}} = 0.22 \pm 0.05 \pm 0.02 \text{ pb}$$


$$(\text{SM } 0.19 - 0.25 \text{ pb})$$




$$e^+e^- \rightarrow ZZ \rightarrow q\bar{q}\nu\bar{\nu}$$

$e^+e^- \rightarrow ZZ \rightarrow q\bar{q}\nu\bar{\nu}$ @ 183 GeV

Run # 671506 Event # 1534 DAQ Time: 03-08-1997 15:18:18




$e^+e^- \rightarrow Z \rightarrow q\bar{q}$ @ 91 GeV





$$e^+e^- \rightarrow ZZ \rightarrow q\bar{q}\nu\bar{\nu}$$




Neural Networks/Likelihoods with:

- Visible mass and energy
- Missing mass and momentum
- Two *vs* three jet topology
- (b-quark content)



$$e^+e^- \rightarrow Z\gamma^* \rightarrow q\bar{q}\nu\bar{\nu}$$

$$Z\gamma^* \rightarrow \nu\bar{\nu}q\bar{q}$$



Defining the signal as $M_{q\bar{q}} < 60$ GeV:



$$\sigma_{e^+e^- \rightarrow Z\gamma^* \rightarrow \nu\bar{\nu}q\bar{q}}^{183 \text{ GeV} - 208 \text{ GeV}} = 0.19 \pm 0.06 \pm 0.02 \text{ pb}$$

(SM 0.13 – 0.16 pb)



$$e^+e^- \rightarrow ZZ \rightarrow q\bar{q}b\bar{b}$$

OPAL preliminary




Neural Networks/Likelihoods with:


- Two *vs* four jet topology
- Kinematic fit to equal mass particles
- b-quark content



$$e^+e^- \rightarrow ZZ \rightarrow l^+l^-l^+l^-$$

	DELPHI	Run: 104731	Evt: 13458												
	Beam: 100.1 GeV	Proc: 6-Aug-1999		Acq	76	82	1	0	1	5	1	0	1	0	1
	DAS: 6-Aug-1999	Scan: 26-Aug-1999		Deact	0	0	0	0	0	0	0	0	0	0	0
	06:52:38	Tan+DST			0	1	0	1	0	1	0	1	0	1	0


$\sqrt{s} = 200 \text{ GeV}$



- Identified leptons
- Invariant masses and opening angles







$$e^+e^- \rightarrow ZZ \rightarrow l^+l^-\nu\bar{\nu}$$





Cross-section measurement


All the experiments fit the discriminating distributions and/or count events, finding a good agreement with the Standard Model predictions:





Cross-section measurement

Combining the results in the NC02 assumption:



$$\sigma_{ZZ}(182.7 \text{ GeV}) = 0.23 \pm 0.08 \pm 0.02 \text{ pb}$$

$$\sigma_{ZZ}(188.7 \text{ GeV}) = 0.70 \pm 0.07 \pm 0.03 \text{ pb}$$

$$\sigma_{ZZ}(191.5 \text{ GeV}) = 0.60 \pm 0.18 \pm 0.04 \text{ pb}$$

$$\sigma_{ZZ}(195.6 \text{ GeV}) = 1.04 \pm 0.12 \pm 0.04 \text{ pb}$$

$$\sigma_{ZZ}(199.6 \text{ GeV}) = 0.98 \pm 0.12 \pm 0.04 \text{ pb}$$

$$\sigma_{ZZ}(201.7 \text{ GeV}) = 0.88 \pm 0.08 \pm 0.04 \text{ pb}$$

$$\sigma_{ZZ}(205.0 \text{ GeV}) = 1.00 \pm 0.15 \pm 0.05 \text{ pb}$$

$$\sigma_{ZZ}(206.8 \text{ GeV}) = 0.70 \pm 0.21 \pm 0.05 \text{ pb}$$




Cross-section measurement

Excellent agreement with the Standard Model predictions


21/07/2000

Summer 00 - Preliminary - Measured $\sigma^{ZZ} / ZZTO$








If we can observe




We are also sensitive to:





Low Scale Gravity



$$\frac{d\sigma}{d\Omega}(s, t) = \frac{d\sigma^{\text{SM}}}{d\Omega}(s, t) + \frac{\lambda}{M_S^4} \alpha_{\text{Interf.}}(s, t) + \frac{\lambda^2}{M_S^8} \beta_{\text{Low Scale Gravity}}(s, t)$$

Extract 95% CL lower limits from a fit to total cross-section/angular information one extracts

	$\lambda = -1$ M_S (TeV)	$\lambda = +1$ M_S (TeV)
L3	1.2	1.2
OPAL	0.6	0.8



- The LEP experiments measure the ZZ cross-section at all the centre-of-mass energies
- Perfect agreement with the Standard Model
- The combined precision of the measurement is 6%
- The measurements of the cross-section into b quarks give confidence to the sensitivity to Higgs boson signals
- No signs of Extra Spatial Dimensions show up in ZZ events, and severe limits are posed on the scale of the so called Low Scale Gravity

I'd like to thank my colleagues for having provided me with their preliminary results, and the LEP Electroweak working group for their combination.