

Measurements of e - γ interactions at LEP

For the LEP collaborations

Carmen Palomares

CIEMAT (Madrid, Spain)



XXXth International Conference on High Energy Physics

July 27 - August 2, 2000, Osaka, Japan

OUTLINE

- ◆ Quasi-real Compton scattering
- ◆ $e^{\pm}\gamma \rightarrow e^{\pm}\gamma$: L3 analysis
- ◆ $e^{\pm}\gamma \rightarrow e^{\pm}\gamma^*/Z$: OPAL and DELPHI analyses
- ◆ Single excited electrons
- ◆ Conclusions

Quasi-real Compton scattering



- Bremsstrahlung:

$$q^2, p^2 \rightarrow 0, \quad e^+, e^- \text{ and } \gamma \text{ undetected.}$$

- Radiative Bhabha scattering:

$$q^2 \rightarrow 0; |p^2| \gg |q^2|.$$

e^+, e^- detected at finite angle, γ along beam direction.

- **Quasi-real Compton scattering:** ($e^\pm \gamma \rightarrow e^\pm \gamma$)

$$p^2 \rightarrow 0; |q^2| \gg |p^2|$$

e^\pm and γ detected ($p_T^\gamma \approx 0$)

the other e^\pm scattered at zero degree.

$$E_{\text{vis}} > E_{\text{beam}}$$

e-γ interactions at LEP

Two processes have been studied:

* $e^\pm \gamma \rightarrow e^\pm \gamma$: L3

Aim: Measurement of $\sigma(e\gamma \rightarrow e\gamma)$.

* $e^\pm \gamma \rightarrow e^\pm Z/\gamma^*$: OPAL, DELPHI

The real photon is replaced by a virtual one or a Z

LEP process $e^+e^- \rightarrow e^+e^-f\bar{f}$

Further processes leading to eeff final state:

- $e^+e^- \rightarrow ZZ \rightarrow e^+e^-f\bar{f}$

- $e^+e^- \rightarrow \gamma\gamma \rightarrow e^+e^-f\bar{f}$

Two channels: $Z/\gamma^* \rightarrow q\bar{q}, \mu^+\mu^-$

Aim: Measurement of $\sigma(e^\pm \gamma \rightarrow e^\pm Z/\gamma^*)$

Signatures:

e^\pm escapes along the beam pipe.

The other e^\pm is observed in the detector together with a γ or two fermions ($Z/\gamma^* \rightarrow f\bar{f}$).

$e^\pm\gamma \rightarrow e^\pm\gamma$: L3 analysis

Quasi-real Compton scattering is studied
at $20 \text{ GeV} < \sqrt{s'} < 185 \text{ GeV}$

Data from 1991-1999, $\mathcal{L} = 634.6 \text{ pb}^{-1}$, $\sqrt{s} \leq 202 \text{ GeV}$

Standard Model predictions:

MC signal ($e^\pm\gamma \rightarrow e^\pm\gamma$): TEEGG $\mathcal{O}(\alpha^4)$

Background: - $e^+e^- \rightarrow \gamma\gamma(\gamma)$.


- $e^+e^- \rightarrow e^+e^-(\gamma)$.

Selection of the signal events

- * γ and e^\pm Identification: signal in electromagnetic calorimeter
- * Polar acceptance for γ -e pair:
 $|\cos\theta| < 0.94$ and $|\cos\theta^*| < 0.80$.
- * To ensure the selection of quasi-real photons:
 $p_T^{e\gamma}/E_{\text{beam}} < 0.15$

RESULTS


$\sqrt{s'}$ – spectra



RESULTS

- ◆ 7335 quiasi-real Compton scattering events
- ◆ Small contribution of virtual photons:
 $\langle P^2/s' \rangle \sim 10^{-3}$

Cross Section



$e^\pm\gamma \rightarrow e^\pm Z/\gamma^*$: OPAL and DELPHI analyses

Electroweak Compton scattering $e^\pm\gamma \rightarrow e^\pm Z/\gamma^*$

Subprocess of the reaction $e^+e^- \rightarrow e^+e^-Z/\gamma^*$; $Z/\gamma^* \rightarrow f\bar{f}$

Signal definition:

OPAL: limits in Lorentz invariant variables

$$|\hat{t}| > 400 \text{ GeV}^2 \quad \hat{t} = (p' - p)^2$$

$$|p^2| < 10 \text{ GeV}^2 \Rightarrow \sqrt{s'} \geq 20.6 \text{ GeV}$$

$$M_{f\bar{f}} > 5 \text{ GeV}$$

DELPHI: topological limits

$$|\cos\theta_e| < 0.985, \quad E_e > 4 \text{ GeV}, \quad M_{f\bar{f}} > 15 \text{ GeV}$$

Standard Model predictions:

MC signal ($e^+e^- \rightarrow e^+e^-Z/\gamma^*$): grc4f, PYTHIA.

Background:

- 4 fermions
- $\gamma\gamma \rightarrow qqee$
- $q\bar{q}(\gamma)$
- 2 fermions

Data from 1997-1999, $183 \text{ GeV} < \sqrt{s} < 202 \text{ GeV}$

$$\underline{e^\pm \gamma \rightarrow e^\pm Z/\gamma^* \rightarrow e^\pm q\bar{q}}$$

Selection

* Preselection:

e^\pm Identification:

Signal in Electromagnetic calorimeter.

Multiplicity

2 jets (Durham algorithm):

OPAL: $M_{jj} > 5 \text{ GeV}$

DELPHI: $M_{jj} > 15 \text{ GeV}$

Kinematic fit: 2 jets + 1 e detec. + 1 e beam pipe
(cut in χ^2)

To ensure signal definition:

OPAL: $\sqrt{s'} = M_{\gamma e} \geq 25 \text{ GeV}$

DELPHI:

$|\hat{t}| > 500 \text{ GeV}^2$

$|\cos \theta_e| < 0.985$

$(\hat{t} \equiv 2 E_{\text{beam}} \cdot E_e (1 + q_e \cdot \cos \theta_e))$

* Selection:

★ Cuts in missing momentum

★ Angular cuts

RESULTS

Low mass region (γ^*ee) $m_{q\bar{q}} < 60$ GeV
 High mass region (Zee) $m_{q\bar{q}} > 60$ GeV

DELPHI


OPAL

$\sqrt{s} = 183$ GeV: First observation

$\sqrt{s} = 189$ GeV



Last Results: Preliminary

Cross section (pb)	γ^*ee	Zee
Measured	$4.6 \pm 0.9 \pm 0.6$	$1.5 \pm 0.3 \pm 0.3$
SM Prediction	3.06 ± 0.04	1.19 ± 0.02



OPAL



Preliminary





DELPHI Preliminary

$\sqrt{s} = 189 \text{ GeV}$



$$\text{DELPHI: } e^{\pm}\gamma \rightarrow e^{\pm}Z/\gamma^* \rightarrow e^{\pm}\mu^+\mu^-$$

- ★ Two tracks to be identified as leptons (μ or e)
- ★ and at least one of them to be a muon.

RESULTS

$\sqrt{s}(\text{GeV})$	N_{expected}	N_{data}	σ (fb)	$\sigma(\text{SM})$ (fb)
$15 < m_{\mu^+\mu^-} < 60 \text{ GeV}$				
182.6	0.5	0	—	112
188.6	1.6	2	154^{+206}_{-129}	112
192-202	2.5	1	—	114
$m_{\mu^+\mu^-} > 60 \text{ GeV}$				
182.6	0.8	1	—	33
188.6	2.4	5	80^{+48}_{-36}	33
192-202	3.8	4	34^{+30}_{-22}	34




Production of single excited electron (L3)



- * The existence of e^* would enhance the number of observed γe events.
- * Since $\sqrt{s'} (\equiv m_{e^*})$ -spectra are in agreement with the SM \implies Upper limit for N_{e^*} from $N_{\gamma e}^{obs}$ and $N_{\gamma e}^{exp,SM}$.

$$\Downarrow \sigma(e\gamma \rightarrow e^* \rightarrow e\gamma) \propto \frac{\lambda^2}{m_{e^*}^2}$$

Upper limit coupling $\lambda (e^*e\gamma)$ as a function of m_{e^*}



CONCLUSIONS

- ◆ γ -e events are identified as Compton scattering of quasi-real photons
 - ▷ The cross-section of this process is measured in the energy range $20 \text{ GeV} < \sqrt{s'} < 185 \text{ GeV}$ and it is in good agreement with the QED expectations.
- ◆ Z/γ^* production in Compton scattering of quasi-real photons has been observed.
 - ▷ The cross-section of the process $\gamma e \rightarrow eZ/\gamma^* \rightarrow eq\bar{q}$ has been calculated separately for Zee-like and γ^*ee -like events, with a cut at a hadronic mass of 60 GeV.
 - ▷ OPAL: The measured cross-section ($\sqrt{s} = 189 \text{ GeV}$) is in agreement with the SM prediction in the Zee region, while in the γ^*ee region an excess with a significance of about 1.4 standard deviations is observed in the data.

- ▷ DELPHI: The measured cross-section ($183 < \sqrt{s} < 202$ GeV) is in agreement with the prediction in both invariant mass regions.
- ▷ The cross-section of the process $\gamma e \rightarrow eZ/\gamma^* \rightarrow e\mu^+\mu^-$ has been measured as well.
- ▷ The results are not conclusive due to low statistics.
- ◆ An upper limit for a hypothetical coupling $e^*e\gamma$ as a function of m_{e^*} is derived from the measurement of Compton scattering.
 - ▷ λ of order of 10^{-1} - 10^{-2} in the region $20 \text{ GeV} < m_{e^*} < 200 \text{ GeV}$.