

Open Charm and Beauty Production at HERA

Felix Sefkow
Universität Zürich

New Results from H1 and ZEUS

Selected Topics:


New tests of perturbative QCD
in ep interactions at $\sqrt{s} = 300\ldots 318 \text{ GeV}$

- Charm production – using $\int \mathcal{L} = 100 \text{ pb}^{-1}$
 - Beauty production – using a vertex detector
-

Charm as a probe of proton structure:
see E. Tzamariudaki's talk in Session PA 03b

Millions of Charm Quarks


- 27,000 $D^* \rightarrow D^0\pi^+ \rightarrow K^-\pi^+\pi^+$ decays




- Production at high x_{Bj} and Q^2
(x, Q^2 : Deep Inelastic Scattering scaling variables)
- Charm spectroscopy

Charm Production

- At low Q^2 : Boson Gluon Fusion





- At higher $Q^2 \gg m_c^2$ charm can be treated as active parton in the proton



In Variable Flavour Number schemes expect deviations from the Boson Gluon Fusion picture

D^* cross sections in DIS


- Compare with NLO QCD (3 Flavour $\overline{\text{MS}}$ scheme)
HVQDIS program (Harris & Smith), Peterson fragmentation
- Use gluon density from scaling violations of proton structure function F_2 (shaded: $m_c = 1.3 - 1.6 \text{ GeV}$)



- Good description up to $Q^2 \simeq 1000 \text{ GeV}^2$ and $x_{Bj} \simeq 0.1$
- Need much more luminosity to detect 1% intrinsic charm contribution at high x_{Bj}

D Meson Spectrum


non-strange mesons



- orbital excitations D_1 and D_2^* established
- radial excitation D^{*J} seen

Neutral P-wave Mesons D_1^0 and D_2^{*0}


- Reconstruction via $D_J^{(*)0} \rightarrow D^{*+}\pi^- + \text{c.c.}$



- Fit invariant mass and π helicity angle
include narrow enhancement at $m(D^*\pi) = 2398$ MeV
- $\frac{D_1^0 \rightarrow D^{*+}\pi^-}{D^{*+}} = 3.40 \pm 0.42^{+0.78\%}_{-0.63\%}$
- $\frac{D_2^{*0} \rightarrow D^{*+}\pi^-}{D^{*+}} = 1.37 \pm 0.40^{+0.96\%}_{-0.33\%}$

Search for radially excited $D^{*''}$

- Reconstruction via $D^{*''+} \rightarrow D^{*+}\pi^+\pi^- + \text{c.c.}$



- $\frac{\langle N_{D^{*''+}} \rangle}{\langle N_{D^{*+}} \rangle} \cdot \text{BR}(D^{*''+} \rightarrow D^{*+}\pi^+\pi^-) < 2.3\% \quad (95\% \text{ C.L.})$
- Compare: $\frac{\langle N_{D^{*''+}} \rangle \text{BR}(D^{*''+} \rightarrow D^{*+}\pi^+\pi^-)}{\sum_{J=1,2} \langle N_{D_J^{(*)0}} \rangle \text{BR}(D_J^{(*)0} \rightarrow D^{*+}\pi^-)}$
 $= 0.49 \pm 0.18 \pm 0.10 \quad (\text{DELPHI})$
 $< 0.21 \quad (95\% \text{ C.L.}) \quad (\text{OPAL})$
 $< 0.16 \quad (95\% \text{ C.L.}) \quad (\text{CLEO})$
- $f(c \rightarrow D^{*''+}) \cdot \text{BR}(D^{*''+} \rightarrow D^{*+}\pi^+\pi^-)$
 $< 0.7\% \quad (95\% \text{ C.L.})$
 $< 1.2\% \quad (95\% \text{ C.L.}) \quad (\text{OPAL})$


Signatures of Beauty

1. Mass

→ rel. transverse momentum p_T^{rel}

2. Lifetime

→ impact parameter δ





Run 186751 Event 18003 Class: 2 4 6 15 20 24 Date 22/07/1999

b cross section at HERA vs. NLO QCD


- Measurements using p_T^{rel}
- ZEUS (e channel)
- H1 (μ channel)

ZEUS Preliminary



b production: impact parameter


- Photoproduction, ≥ 2 jets $E_T > 5$ GeV,
 $\geq 1 \mu$ in 2 layer silicon vertex detector



- Likelihood fit: $f_b = 26 \pm 5\%$
- Cross section in kinematic range
 $Q^2 < 1 \text{ GeV}^2$, $0.1 < y < 0.8$, $p_T(\mu) > 2 \text{ GeV}$, $35^\circ < \theta(\mu) < 130^\circ$
- $\sigma_{vis}(ep \rightarrow b\bar{b}X \rightarrow \mu X) =$
 $[159 \pm 30 (\text{stat.}) \pm 29 (\text{syst.})] \text{ pb}$
- Confirms published H1 result $(176 \pm 16^{+27}_{-17}) \text{ pb}$

b production: combine impact parameter δ and p_T^{rel}

- High b purity regions



- Combined likelihood fit in (δ, p_T^{rel}) plane:
 $\sigma_{vis}(ep \rightarrow b\bar{b}X \rightarrow \mu X) = [160 \pm 16 (\text{stat.}) \pm 29 (\text{syst.})] \text{ pb}$
- Average with publ. H1 result: $\sigma_{vis} = (170 \pm 25) \text{ pb}$
- NLO QCD (3 Flavour): $\sigma_{vis} = (104 \pm 17) \text{ pb}$

Open Beauty Production:

- First b cross section measurement confirmed with lifetime based method
- Excess over NLO QCD established

Conclusion:

With plenty of Charm
and long-lived Beauty:

HERA

takes another step forward in precision
to probe the strong interactions