

# Excited leptons and leptoquarks at LEP

**M.A. Falagan**

**CIEMAT**

**XXXth International Conference on High Energy  
Physics**


**July 27 - August 2, 2000, Osaka, Japan**

- Excited leptons
- Leptoquarks
- Summary


- Excited fermions are expected in composite models. These models could explain the family problem and make the fermion masses calculable parameters.
- A well studied model, assumes spin 1/2, and isospin doublets with left and right handed components.

$$L^* = \begin{pmatrix} \nu^* \\ \ell^* \end{pmatrix}_L + \begin{pmatrix} \nu^* \\ \ell^* \end{pmatrix}_R$$

- They can be produced in pairs or singly.



- The cross section for pair production is determined by the charge and the isospin. It depends only on  $\sqrt{s}$  and  $m_*$ .
- The cross section for single production depends on the effective couplings  $f$  and  $f'$  and  $m_*$ .
- An excited lepton decays immediately into a standard lepton plus a gauge boson ( $\gamma$ , W or Z).



Pair produced excited leptons have been extensively searched for in LEP, in many different signatures, corresponding to the dominant decay modes.

$\sqrt{s} \sim 200$ GeV	L3		OPAL	
Signal	Data	SM	Data	SM
$e^*e^* \rightarrow e\gamma e\gamma$	1	0.8	2	1.4
$\mu^*\mu^* \rightarrow \mu\gamma\mu\gamma$	1	0.5	0	0.7
$\tau^*\tau^* \rightarrow \tau\gamma\tau\gamma$	0	0.2	3	2.6

Data and SM expectation always agrees  $\implies$

lower mass limits are set


Excited Lepton	95% CL Mass Limit (GeV)			
	$f = f'$		$f = -f'$	Coup. Indep.
	L3	OPAL	L3	L3
$e^*$	100.1	100.1	96.2	96.0
$\mu^*$	100.3	100.1	96.2	96.2
$\tau^*$	99.9	100.0	96.2	94.9
$\nu_e^*$	99.3	99.5	99.5	98.5
$\nu_\mu^*$	99.4	99.5	99.5	98.5
$\nu_\tau^*$	93.9	91.9	99.4	92.7

# Candidate to $e^*e^*$ production. L3 $\sqrt{s} = 200$ GeV


$m(e_1, \gamma_2) = 60$  GeV

$m(e_2, \gamma_1) = 59$  GeV

Run # 772501 Event # 3103 Total Energy : 185 GeV







# Experimental distributions from the $l\bar{l}\gamma$ selection







OPAL Preliminary

OPAL Preliminary



Experimental distributions from the  $\ell\nu W$  selection





## FROM SINGLE PRODUCTION SEARCHES

No excess in data is seen


Limits are derived on  $f_0/\Lambda = \sqrt{(f^2 + f'^2)}/2/\Lambda$  independently of the coupling assumption  $f/f'$

## OPAL



(From data at  $\sqrt{s} = 161 - 183$  GeV)


Including data at higher energies (192 – 202 GeV) and considering two particular scenarios ( $f = f'$  and  $f = -f'$ ) improved limits are derived on the coupling constant  $f/\Lambda$



Similar limits are derived by DELPHI and OPAL




The search for excited electrons is extended above the kinematical limit for single production ( $m_* = \sqrt{s}$ )



Assuming the coupling strength of the  $e^*e\gamma$  vertex equal to the standard  $ee\gamma$  vertex, a mass limit to  $e^*$  is derived from the differential cross section for the process  $e^+e^- \rightarrow \gamma\gamma(\gamma)$




OPAL  $\sigma(e^+e^- \rightarrow \gamma\gamma(\gamma)) \implies m_* > 316 \text{ GeV}$  at 95% C.L.



Leptoquarks are new bosonic fields which mediate interaction between quark and leptons. They carry baryon and lepton number and therefore decay into a pair lepton - quark.


They have color, electric charge and weak isospin, and therefore couple to the standard gauge bosons.

Assuming dimensionless interactions with Standard Model fermions and gauge invariance, there could be 9 scalar states **S** and 9 vector states **V**, grouped in two singlets, two doublets and a triplet of isospin.



Pair produced leptoquarks have been searched for in the Opal experiment at  $\sqrt{s} = 189 - 202$  GeV in all their decay channels

**OPAL preliminary Selection: LQ LQ => jet jet nu nu**



No excess in Data  $\implies$  Mass limits are set on scalar leptoquarks

95% C.L. Mass limits (GeV)

LQ	$Q_{em}$	$Br(\text{LQ} \rightarrow \ell q)$	1 <sup>st</sup> gen.	2 <sup>nd</sup> gen.	3 <sup>rd</sup> gen.
$S_0$	$-1/3$	[0.5, 1.]	44.2*	44.2*	41.4*
$\tilde{S}_0$	$-4/3$	1.	94.7	96.4	95.2
$S_{1/2}$	$-2/3$	[0., 1.]	88.5	88.9	89.4
	$-5/3$	1.	95.6	97.0	96.1
$\tilde{S}_{1/2}$	$1/3$	0.	83.3	83.3	83.3
	$-2/3$	1.	90.5	94.6	93.1
$S_1$	$2/3$	0.	91.7	91.7	91.7
	$-1/3$	0.5	44.2*	44.2*	43.2*
	$-4/3$	1.	96.0	97.4	96.5

(\*) Limits from LEP1

Singly produced leptoquarks have been searched for in the Opal (189 GeV) and Delphi (192 – 202 GeV) experiments.

Signal	OPAL		DELPHI	
	Data	SM	Data	SM
$e^+e^- \rightarrow e q \text{ LQ} \rightarrow e q e q$	21	22	6	6
$e^+e^- \rightarrow e q \text{ LQ} \rightarrow e q \nu q$	7	9	5	4

No excess in Data  $\implies$

assuming coupling parameter to fermions  $\lambda = \sqrt{4\pi\alpha_{em}}$

mass limits are set on first generation scalar and vector leptoquarks

95% C.L. Mass limits (GeV) on scalar LQ

Leptoquark	$Q_{em}$	$Br(\text{LQ} \rightarrow \ell q)$	OPAL	DELPHI
$S_0$	-1/3	0.5	158	180
		1.	163	180
$\tilde{S}_0$	-4/3	1.	149	158
$S_{1/2}$	-2/3 -5/3	1.	121	155
		1.	164	180
$\tilde{S}_{1/2}$	1/3 -2/3	0.		
		1.	121	155
$S_1$	2/3 -1/3 -4/3	0.		
		0.5	158	180
		1.	156	158


- Lower mass limits are also set on vector leptoquarks

95% C.L. Mass limits (GeV) on vector LQ

Leptoquark	$Q_{em}$	$Br(\text{LQ} \rightarrow \ell q)$	OPAL	DELPHI
$V_{1/2}$	$-1/3$	1.	176	188
	$-4/3$	1.	152	168
$\tilde{V}_{1/2}$	$2/3$	0.		
	$-1/3$	1.	176	188
$V_0$	$-2/3$	0.5	149	171
		1.	151	170
$\tilde{V}_0$	$-5/3$	1.	177	185
$V_1$	$1/3$	0.		
	$-2/3$	0.5	149	171
	$-5/3$	1.	182	185

The study of  $\sigma(e^+e^-) \rightarrow q\bar{q}$  extends the sensitivity to leptoquarks above the kinematical limit for single production ( $m_{LQ} = \sqrt{s}$ )

The  $t$ -channel LQ exchange diagrams would be an additional contribution to hadron production in LEP



Assuming coupling parameter to fermions  $\lambda = \sqrt{4\pi\alpha_{em}}$   
 mass limits are set on scalar and vector leptoquarks

Leptoquark	L3	ALEPH
$S_0(L/R) \rightarrow eu$	413 / 322	380 / 56
$\tilde{S}_0(R) \rightarrow ed$	84	128
$S_{1/2}(L/R) \rightarrow e\bar{u} / e\bar{u}, e\bar{d}$	64 / 117	120 / 99
$S_1(L) \rightarrow eu, ed$	208	319
$V_0(L/R) \rightarrow e\bar{d}$	584 / 136	618 / 137
$\tilde{V}_0(R) \rightarrow e\bar{u}$	288	331
$V_{1/2}(L/R) \rightarrow ed / eu, ed$	202 / 183	144 / 169
$\tilde{V}_{1/2}(L) \rightarrow eu$	145	105
$V_1(L) \rightarrow e\bar{u}, e\bar{d}$	394	515

- Excited leptons and leptoquarks have been searched for in pair production, single production and indirectly at LEP.
- No evidence for their existence has been found.
- From pair production searches, lower limits on the mass of excited leptons and leptoquarks have been set at the 95% confidence level.
- From single production and indirect searches upper limits to the couplings as a function of the mass have been derived.
- With the current LEP run at  $\sqrt{s} = 208$  GeV the range of discovery potential has been extended around 3 GeV for pair production and 6 GeV for single production. In case no signal appears, limits will improved.