

R-parity violation searches at LEP

Ivor Fleck

University of Freiburg

July 28, 2000

PA-11

- R-parity violating superpotential
- indirect limits on couplings
- experimental results
 - trilinear terms
 - * gauginos
 - * sleptons
 - * sneutrinos
 - * squarks
 - bilinear terms
- conclusions and outlook

Introduction of R-parity

new discrete multiplicative symmetry in SUSY models

$$R_p = (-1)^{2S+3B+L}$$

S: spin, B: baryon number, L: lepton number

$$\begin{aligned} R_p &= \mathbf{1} && \text{for SM particles} \\ R_p &= \mathbf{-1} && \text{for SUSY particles} \end{aligned}$$

	R_p	\cancel{R}_p
SUSY particles produced in	pairs	pairs or singly
the LSP	is stable	decays
experimental signature	\cancel{E}_T	E_T


Exclusion limits under assumption of R_p -conservation are not valid under R_p -violation.

The R-parity Violating Superpotential



$$W = \lambda_{ijk} L_L^i L_L^j \bar{E}_R^k + \lambda'_{ijk} L_L^i Q_L^j \bar{D}_R^k + \lambda''_{ijk} \bar{U}_R^i \bar{D}_R^j \bar{D}_R^k + \epsilon_i L_L^i H_u$$

with

$L_L(Q_L)$:	lepton (quark) doublet superfield
E_R :	electron singlet superfield
$D_R(U_R)$:	down- (up-) quark type singlet superfield
H_u :	Higgs field coupling to up-quarks
$\lambda, \lambda', \lambda''$:	Yukawa couplings
ϵ :	effective coupling
i, j, k :	generation indices



L violation



B violation

Due to symmetrie $i < j$ for λ_{ijk}

$j < k$ for λ''_{ijk}


→ 48 (9 + 27 + 9 + 3) independent couplings

Limits for R-parity Violation

R_p -violation leads to several measurable effects e.g.

- lepton universality violation
- fast proton decay

Non observation of these effects sets limits on the size of the λ -couplings



λ	λ'			λ''
(121) 0.05	(111) 0.0005	(211) 0.06	(311) 0.1	(121) 10^{-15}
(122) 0.05	(112) 0.02	(212) 0.06	(312) 0.1	(131) 10^{-4}
(123) 0.05	(113) 0.02	(213) 0.06	(313) 0.1	(132) 1.23
(131) 0.06	(121) 0.04	(221) 0.18	(321) 0.52	(221) 1.23
(132) 0.06	(122) 0.04	(222) 0.21	(322) 0.52	(231) 0.5
(133) 0.006	(123) 0.04	(223) 0.21	(323) 0.52	(232) 0.5
(231) 0.07	(131) 0.02	(231) 0.18	(331) 0.45	(321) 0.5
(232) 0.07	(132) 0.3	(232) 0.56	(332) 0.45	(331) 0.42
(233) 0.07	(133) 0.001	(233) 0.15	(333) 0.45	(332) 0.42

2σ limits for sparticle mass of 100 GeV

red numbers indicate couplings excluded at 2σ from a global fit to hadronic Z decays

More stringent limits on product of two λ -couplings


– non observation of proton decay:

$$\lambda'_{11k} \times \lambda''_{11k} < 10^{-24}$$

→ assume exactly one λ -coupling is different from 0

Decay Width of Gauginos

Decay width of photino like $\tilde{\chi}_1^0$ from pair-production of $\tilde{\chi}_1^0$



$$\Gamma = \lambda^2 \frac{\alpha}{128\pi^2} \frac{(m_{\tilde{\chi}_1^0})^5}{(m_{\tilde{f}})^4}$$

$m_{\tilde{f}}$: virtual sfermion mass (set to 100 GeV)

→ sensitive to λ -values of $\mathcal{O}(10^{-5})$



decay width depends strongly on $\tilde{\chi}_1^0$ matrix

Gauginos

two different decay modes:

R_p violating =
direct

R_p conserving + R_p violating =
indirect



final states from pair-production:

	direct	indirect
gauginos	6 fermions	10 fermions

direct: λ

$$\tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow 4 l + \cancel{E}$$

$$\tilde{\chi}_1^+ \tilde{\chi}_1^- \rightarrow 2 l + \cancel{E}$$

$$\tilde{\chi}_1^+ \tilde{\chi}_1^- \rightarrow 4 l + \cancel{E}$$


$$\tilde{\chi}_1^+ \tilde{\chi}_1^- \rightarrow 6 l$$

indirect: λ



$$\tilde{\chi}_1^+ \tilde{\chi}_1^- \rightarrow W W + 4 l + \cancel{E}$$

limits shown for λ -coupling resulting in worst cross-section limit


OPAL



Limits from Gaugino Searches



exclusion up to kinematic limit of $\tilde{\chi}_1^\pm$ pair production
 extended sensitivity due to $\tilde{\chi}_1^0$ pair production



exclusions:

$$M(\tilde{\chi}_1^0) > 35 \text{ GeV}$$

$$M(\tilde{\chi}_2^0) > 68 \text{ GeV}$$

$$M(\tilde{\chi}_1^\pm) > 99 \text{ GeV}$$


independent of

$\tan\beta, m_0, M_2, \mu$


$\lambda, \lambda', \lambda''$

Sleptons

direct




indirect



final states:

	direct	indirect
λ	$2\ell + \cancel{E}$	$6\ell + \cancel{E}$
λ'	4 jets	$(2+x)\ell + 4 \text{ jets } (x=0,1,2)$
λ''	-	$2\ell + 6 \text{ jets}$




upper cross-section limits for $2\ell + \cancel{E}$ final states




Limits from Slepton Searches

Exclusions from slepton searches: (e.g. λ)

OPAL Preliminary



Exclusions from $\tilde{\chi}_1^\pm$ searches:
(t-channel exchange of selectron)




Sneutrinos

final states:

	direct	indirect
λ	4ℓ	$4 \ell + \cancel{E}$
λ'	4 jets	$x \ell + 4 \text{ jets} + \cancel{E} \quad (x=0,1,2)$
λ''	-	$6 \text{ jets} + \cancel{E}$



decays via λ :

sneutrino limits from different centre-of-mass energies





Sneutrino limits

limits from decays via λ' :



limits from decays via λ'' :




Single Sneutrino Production

DELPHI preliminary

direct decay:




2 fermion final states:





$$\begin{aligned} \mu^+ \mu^- : & \quad \lambda_{131} = \lambda_{232} \\ \tau^+ \tau^- : & \quad \lambda_{121} = \lambda_{323} \end{aligned}$$

sensitivity for $\lambda > \mathcal{O}(10^{-2})$

indirect decay:



ALEPH Preliminary



sensitivity for
 $\lambda > \mathcal{O}(10^{-3})$
for masses
up to \sqrt{s}

Squark Decays

mixing of left- and right handed scalar squarks

$$\tilde{t}_1 = \cos \theta_{\tilde{t}} \tilde{t}_L + \sin \theta_{\tilde{t}} \tilde{t}_R$$

For $\theta_{\tilde{t}} = 0.98(1.17)$ the \tilde{t}_1 (\tilde{b}_1) decouples from Z^0 and the production cross section becomes minimal.


final states:

	direct	indirect
λ	—	2 jets + 4 ℓ + E
λ'	2 jets + 2 (1,0) ℓ	6 jets + 2 (1,0) ℓ
λ''	4 jets	8 jets


indirect decay via λ :

$$\tilde{q} \rightarrow q + \tilde{\chi}_1^0$$

ALEPH Preliminary




ALEPH Preliminary



Squark Decays via λ'


for $\lambda' > 10^{-4}$ direct \tilde{t}_1 decay dominant : $\tilde{t}_1 \rightarrow q + \ell$
 2 jet + 2 lepton final states:



$\tilde{t}_1 \rightarrow q + e$ and $\tilde{t}_1 \rightarrow q + \mu$ excluded
 from CDF and D0 for $M(\tilde{t}_1) < 200$ GeV
 but $\tilde{t}_1 \rightarrow q + \tau$ not excluded


LEP: lower mass limits for
 any mixing angle
 $M(\tilde{t}_1) > 91$ GeV

indirect decay via λ' :



Single Squark production

single production:




\tilde{q} masses up to \sqrt{s} accessible for $\lambda' > 10^{-1}$

t-channel exchange: λ'_{ijk}

final state: 2 jets

\tilde{q} masses even above \sqrt{s} accessible for $\lambda' > 10^{-1}$



Spontaneous R-parity violation



spontaneous breaking of Supersymmetry:

- massless Majoron (J)
- R_p violating superpotential $\epsilon_i L_i H_u$
- mixing between lepton and gaugino sector (3. generation)
- leading to neutrino masses

- additional decays:

$$\tilde{\chi}_1^\pm \rightarrow \tau^\pm J$$

signal signature:
two acoplanar taus



at $\sqrt{s} = 183$ and 189 GeV

15 events observed with

15.8 events expected

$$m(\tilde{\chi}_1^\pm) > 94.4 \text{ GeV}$$

for $m(\tilde{\nu}) > 300$ GeV

Conclusions and Outlook

- Searches for R_p -violating production and decay of SUSY particles are complementary to searches with R_p -conservation
- Searching for both scenarios has extended discovery potential
- in case of R_p -violation sparticles masses up to \sqrt{s} can be probed
- Under the assumption of R_p -violation searches have been performed at LEP for $\sqrt{s} \leq 202$ GeV gauginos, sleptons, sneutrinos and squarks for all Yukawa couplings λ , λ' , λ'' and for bilinear terms ϵ
- No evidence for SUSY with R_p -violation so far at LEP
- neutrino masses can be explained with R_p -violation
- results from this years data will be presented on Saturday
- R_p -violating scenarios have to be considered at future colliders like LHC or NLC