


**Inclusive Particle Production
in Two-Photon Collisions
at LEP**

Christoph Schäfer
CERN

ICHEP 2000, Osaka

July 29th, 2000





Two-Photon Collision Process:




Photons don't interact with each other
(\Rightarrow Abelian structure of QED!),
but can interact within a “**hadronic blob**”

- Look for π^0 , K_S^0 or charged hadrons (h^\pm) in the detector
- Measure their transverse momentum distribution
- Check QCD predictions

Hadronic Two-Photon Collision Process:



- Electron and positron escape undetected along the beam pipe
→ **quasi real photons**
- Relative contribution of Feynman diagrams depends on p_T -range



Two-Photon cross section dominant at LEP

⇒ Check of perturbative QCD

L3: (π^0 , K_S^0)

Integrated Luminosity: 410 pb⁻¹

at 189 < \sqrt{s} < 202 GeV

(2 million events)

All data preliminary

Opal: (h^\pm , K_S^0)

Integrated Luminosity: 20 pb⁻¹

at 161 < \sqrt{s} < 172 GeV

(58 000 events)

(Eur. Phys. J. C 6 (1999) 253)


Monte Carlo Simulation used:

PHOJET: R. Engel and J. Ranft

(Phys. Rev. D54, 4246, 1996, V1.05c)

PYTHIA: T. Sjöstrand


(CPC 82, 74, 1994, V5.718)



Typical $\gamma\gamma \rightarrow \text{hadrons}$ selection:

- low visible energy
- at least few particles in the detector from the interaction point
- small energy deposit in the luminosity monitor (anti tag)

$$e^+e^- \rightarrow e^+e^- + h^\pm + X$$




At low p_T : (< 1 GeV)

$\gamma\gamma$ behaves like soft hadron-hadron and photon-hadron collisions

At high p_T : (> 1 GeV)

Hard QCD and direct QED processes become dominant

Look for two photons from the π^0 -decay



Transverse momentum range of the π^0 :



$$0.2 \text{ GeV} < p_T < 7.5 \text{ GeV}$$

Pseudo-Rapidity: $\eta = -\ln(\tan\theta/2)$

$$|\eta| < 0.5 \quad (\text{Barrel-Region})$$

Differential Cross Sections:

$$e^+e^- \rightarrow e^+e^- + \pi^0 + X$$



Exponential fit for $p_T < 1 \text{ GeV}$:


$$\langle p_T \rangle = 239 \pm 3 \text{ MeV}$$

Compatible with pp- and $p\gamma$ -collisions

NLO QCD: J. Binnewies et al. (priv. com.)

Phys. Rev. D56 (1996) 6110

Look for two charged Pions from the K_S^0 -decay
with secondary vertex




Invariant Mass of the $\pi^+\pi^-$ -system

Pseudo-Rapidity $|\eta| < 1.5$ (Barrel + Endcap)

Differential Cross Sections:

$$e^+e^- \rightarrow e^+e^- + K_S^0 + X$$




Exponential fit for $p_T < 2$ GeV:


$$\langle p_T \rangle = 357 \pm 8 \text{ MeV}$$

Kink at $p_T \approx 3$ GeV due to charm production

Hadrons/Pions



Kaons



Good agreement between L3 and OPAL

- Clean π^0 and K_S^0 signal in $\gamma\gamma$ -collisions
- First study of π^0 and K_S^0 differential cross sections at LEP
- Measurements from L3 and OPAL are consistent
- At low p_T , exponential behaviour in both channels; typical for soft $pp/p\gamma$ interactions
- At high p_T , evidence for contributions of hard QCD and direct QED processes