


Year 2000 update for L3

Simonetta Gentile

CERN EP, Università "La Sapienza"(Roma)



XXXth International Conference on High Energy Physics

27 July - 2 August, Osaka, Japan.



Outline

PRELIMINARY results on 2000 data by L3 experiment on:



- \sqrt{s} up to 205.5 GeV data, $\int \mathcal{L} dt \approx 62 \text{ pb}^{-1}$
- \sqrt{s} up to 207.5 GeV data, $\int \mathcal{L} dt \approx 32 \text{ pb}^{-1}$
- $\sqrt{s} \geq 207.5$ GeV data, $\int \mathcal{L} dt \approx 1.8 \text{ pb}^{-1}$

- Search for Higgs bosons.
 - Standard Model Higgs
 - MSSM Higgs
 - Charged Higgs
 - Invisible Higgs, fermiophobic Higgs
- Search for SUSY particles.
 - Scalar Leptons, Charginos and Neutralinos
 - Scalar Top and Scalar Bottom Quarks
 - Single and Multi-Photons events with missing energy
- Search for exotica and new physics.
 - Heavy Neutral and Charged Leptons
 - Excited Leptons




Four jet channel $HZ \rightarrow qqqq$

Mass, b tag and discriminating variable distribution






$HZ \rightarrow \text{qqqq}$ candidate ($\sqrt{s} = 206.6$ GeV)



$m_H = 109.9$ GeV, event b tag = 2.6



Combined mass plot of this year data after cut on b tag




observed / expected events

Channel	Data	BG	Signal 107 GeV	$m_H \geq 100$ GeV		
				Data	BG	Signal 107 GeV
qqqq	9	9.9	2.8	4	2.9	2.0
qq $\nu\nu$	5	4.6	2.0	1	2.8	1.9
qq $\ell\ell$	0	1.5	0.5	0	0.2	0.4
Sum	14	16.0	5.3	5	5.9	4.3




No signal observed
 → limits set at 95% CL



Observed: $m_H > 107.6 \text{ GeV}$
 median expected: $m_H > 108.5 \text{ GeV}$



Four jet channel $hA \rightarrow bbbb$



			$m_h + m_A \geq 160 \text{ GeV}$		
Data	BG	Signal 90 GeV	Data	BG	Signal 90 GeV
9	7.9	2.9	3	3.5	2.1

MSSM high $\tan \beta$ limit:


$m_h > 87 \text{ GeV}$ (expected 87 GeV)

$m_A > 87 \text{ GeV}$ (expected 87 GeV)




Searches for Charged Higgs Bosons $H^\pm \rightarrow cs, \tau\nu$

Higgs candidate masses
(this year data, $\int \mathcal{L} dt \approx 92.1 \text{ pb}^{-1}$)





Invisible Higgs excluded regions in the plane M_H vs. $\text{BR}(H \rightarrow \text{invisible})$




$$R_{\text{inv}} = \frac{\text{BR} (h \rightarrow \text{inv}) \times \sigma(e^+e^- \rightarrow h \bar{Z})}{\sigma(e^+e^- \rightarrow H_{\text{SM}} \bar{Z})}$$




Searches for Fermiophobic Higgs Bosons

New Analysis: $H \rightarrow WW^*$


Production mechanism



reconstructed mass



Limits on Cross section \times Br($H \rightarrow WW^*$)





- At $\sqrt{s} > 202$ GeV searches in the following channels are performed:

- $\tilde{\chi}_1^\pm, \tilde{\chi}_1^0$
- $\tilde{e}, \tilde{\mu}, \tilde{\tau}, \tilde{t}, \tilde{b}$
- $\gamma + \text{missing energy}, \gamma\gamma + \text{missing energy}$:

- Basic assumptions:

R-parity conservation: pair production of S-particles

and Lightest S-particle (LSP) $\tilde{\chi}_1^0$ stable and invisible

- Sleptons: $e^+ e^- \rightarrow \tilde{l}\tilde{l} \rightarrow l l \tilde{\chi}_1^0 \tilde{\chi}_1^0$
2 leptons, E_{miss}
- Squarks: $e^+ e^- \rightarrow \tilde{q}\tilde{q} \rightarrow q\bar{q} \tilde{\chi}_1^0 \tilde{\chi}_1^0$
2 jets, E_{miss}
- Charginos: $e^+ e^- \rightarrow \tilde{\chi}_1^+ \tilde{\chi}_1^- \rightarrow W^* W^* \tilde{\chi}_1^0 \tilde{\chi}_1^0$
 E_{miss} and 4 jets, or 2 jets and 1 lepton, or 2 leptons

$$\Delta M = (M_{Sparticle} - M_{LSP})$$

Most important parameter for all analyses



- 30 selections for Charginos, Neutralinos, Sleptons

Neutralinos

$\sqrt{s} = 202 - 208 \text{ GeV}, \int \mathcal{L} dt = 56.7 \text{ pb}^{-1}$

$\sqrt{s} > 205.5 \text{ GeV}, \int \mathcal{L} dt = 14.0 \text{ pb}^{-1}$ (high energy sample)

	Low ΔM		Medium ΔM		High ΔM		.OR.	
	N_{data}	N_{exp}	N_{data}	N_{exp}	N_{data}	N_{exp}	N_{data}	N_{exp}
$\tilde{\chi}_2^0$	11	13.3	1	3.5	0	1.2	12	17.4
$\tilde{\chi}_2^0$	4	3.3	0	0.9	0	0.3	4	4.3

$$N_{\tilde{\chi}_2^0}^{\text{Exp}} \sim 28 \quad N_{\tilde{\chi}_2^0}^{\text{Exp}} \sim 7$$

Charginos

$\sqrt{s} = 202 - 208 \text{ GeV}, \int \mathcal{L} dt = 84.3 \text{ pb}^{-1}$

$\sqrt{s} > 205.5 \text{ GeV}, \int \mathcal{L} dt = 28.0 \text{ pb}^{-1}$ (high energy sample)

	Low ΔM		Medium ΔM		High ΔM		.OR.	
	N_{data}	N_{exp}	N_{data}	N_{exp}	N_{data}	N_{exp}	N_{data}	N_{exp}
$\tilde{\chi}_1^\pm$	20	16.7	17	10.5	9	13.8	44	37.4
$\tilde{\chi}_1^\pm$	7	5.4	3	3.4	3	4.6	13	12.4


$$N_{\tilde{\chi}_1^\pm}^{\text{Exp}} \sim 42 \quad N_{\tilde{\chi}_1^\pm}^{\text{Exp}} \sim 14$$

No excess respect to SM predictions

- Enough $\int \mathcal{L} dt$ for $\tilde{\chi}_1^\pm$ exclusions



- Setting model independent chargino mass limits is now possible by combining: standard missing energy search with ISR analysis and highly ionising track analysis.
- New results from ISR analysis. Assuming no signal in 1998–1999 past years data used as precise background prediction.



selection	low ΔM $\Delta M \sim 3\text{GeV}$	very low ΔM $\Delta M \sim 1\text{GeV}$	ultra low ΔM $\Delta M \sim 300\text{MeV}$	OR
1998 (176 pb)	6	1	1	8
1999 (232 pb)	8	4	2	9
2000 (76.5 pb)	3	1	3	5
98-99 scaled	2.6 ± 0.7	0.9 ± 0.4	0.6 ± 0.3	3.2 ± 0.8



Standard Susy Searches


$\sqrt{s} \approx 205 \text{ GeV}, \int \mathcal{L} dt = 84.6 \text{ pb}^{-1}$

	Low ΔM		Medium ΔM		High ΔM		.OR.	
	N_{data}	N_{exp}	N_{data}	N_{exp}	N_{data}	N_{exp}	N_{data}	N_{exp}
\tilde{e}	9	9.9	0	4.3	5	9.2	14	20.4
$\tilde{\mu}$	17	13.6	6	5.4	14	12.8	33	27.5
$\tilde{\tau}$	76	66.4	21	21.6	17	12.5	89	76.6

$$N_{\tilde{e}, \tilde{\mu}, \tilde{\tau}}^{\text{exp}} \sim 4 - 10$$

smuons




staus





S-lepton mass limits

- New s-lepton mass limits including all data from 189 to 208 GeV





At $\sqrt{s} \sim 205\text{GeV}$ $\int \mathcal{L} dt = 49.3\text{pb}^{-1}$

- Scalar top

$$e^+ e^- \rightarrow \tilde{t}_1 \bar{\tilde{t}}_1 \rightarrow c\tilde{\chi}_1^0 \bar{c}\tilde{\chi}_1^0 \implies 2\text{jets} + E_{\text{miss}}$$

$$e^+ e^- \rightarrow \tilde{t}_1 \bar{\tilde{t}}_1 \rightarrow b\ell^+ \bar{\nu} \bar{b}\ell^- \bar{\nu} \implies 2\text{b-jets} + 2\ell + E_{\text{miss}}$$

- Scalar bottom

$$e^+ e^- \rightarrow \tilde{b}_1 \bar{\tilde{b}}_1 \rightarrow b\tilde{\chi}_1^0 \bar{b}\tilde{\chi}_1^0 \implies 2\text{b-jets} + E_{\text{miss}}$$

Channel	Vlow ΔM		Low ΔM		Medium ΔM		High ΔM		OR	
	Data	N_{MC}	Data	N_{MC}	Data	N_{MC}	Data	N_{MC}	Data	N_{MC}
$\tilde{t}_1 \rightarrow c\tilde{\chi}_1^0$	21	14.2	0	2.12	3	2.00	4	3.03	28	21.1
$\tilde{t}_1 \rightarrow b\ell\bar{\nu}$	10	8.11	5	2.52	1	0.20	-	-	16	10.60
$\tilde{t}_1 \rightarrow b\tau\bar{\nu}$	4	4.98	2	4.08	1	0.91	-	-	7	9.39
$\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0$	13	9.93	1	1.138	0	0.34	0	0.33	14	11.74

No significant excess with respect to SM





$\gamma(\gamma) + \text{missing energy}$

Standard Model Process $e^+e^- \rightarrow \nu\bar{\nu}\gamma(\gamma)$

In the GMSB scenario, with \tilde{G} LSP,
the possible decay channels are :

- $e^+e^- \rightarrow \tilde{\chi}_1^0 \tilde{G} \rightarrow \tilde{G}\tilde{G}\gamma$
- $e^+e^- \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow \tilde{G}\tilde{G}\gamma\gamma$

$$\sqrt{s} > 202 \text{ GeV} \quad (\theta \geq 14 \text{ deg})$$



Rule out SUSY interpretation
of CDF $ee\gamma\gamma$ event



New Heavy Leptons


Search for new Unstable Neutral Heavy Leptons. $\mathcal{L} = 96 \text{ pb}^{-1}$

Channel	Data	Background
$e^+e^- \rightarrow L^0L^0 \rightarrow eWeW$	2	2.1
$e^+e^- \rightarrow L^0L^0 \rightarrow \mu W\mu W$	1	0.6
$e^+e^- \rightarrow L^0L^0 \rightarrow \tau W\tau W$	20	26.5

No signal - good agreement between the number of candidates and expected background

Search for Stable Charged Heavy Leptons

103.4 GeV and above, $L=23\text{pb}^{-1}$



Two highly ionising tracks are searched for
No signal \implies Mass $> 102.4 \text{ GeV}$







Search for Pair Production of Excited Leptons. $\mathcal{L} = 90 \text{ pb}^{-1}$

Signal	Data	Bkg	Signal	Data	Bkg
$e^*e^* \rightarrow ee\gamma\gamma$	0	0.3	$\nu_e^*\nu_e^* \rightarrow eeWW$	0	0.1
$\mu^*\mu^* \rightarrow \mu\mu\gamma\gamma$	0	0.2	$\nu_\mu^*\nu_\mu^* \rightarrow \mu\mu WW$	0	0.3
$\tau^*\tau^* \rightarrow \tau\tau\gamma\gamma$	0	0.1	$\nu_\tau^*\nu_\tau^* \rightarrow \tau\tau WW$	1311	1255
$\ell^*\ell^* \rightarrow \nu\nu WW$	1311	1255	$\nu^*\nu^* \rightarrow \nu\nu\gamma\gamma$	1	0.7
$e^*e^* \rightarrow e\gamma\nu W$	3	2.6	$\nu_e^*\nu_e^* \rightarrow \nu\gamma eW$		
$\mu^*\mu^* \rightarrow \mu\gamma\nu W$	1	1.3	$\nu_\mu^*\nu_\mu^* \rightarrow \nu\gamma\mu W$	2	2.3
$\tau^*\tau^* \rightarrow \tau\gamma\nu W$	15	13	$\nu_\tau^*\nu_\tau^* \rightarrow \nu\gamma\tau W$		

No signal \implies Mass limits are set

Excited Lepton Coupling	95% CL Mass Limit (GeV)		
	$f = f'$	$f = -f'$	Coup. Indep.
e^*	102.5	95.9	95.7
μ^*	102.5	95.9	95.9
τ^*	102.4	95.9	94.9
ν_e^*	101.5	102.1	101.3
ν_μ^*	101.8	102.1	101.3
ν_τ^*	93.5	101.9	92.7

From single production searches, limits are derived on the coupling constant $|f|/\Lambda$





Conclusions

21

Preliminary results on 2000 data by L3 experiment has been presented based on data at $\sqrt{s} = 202 - 208 \text{ GeV}$, with $\int \mathcal{L} dt \approx 90 \text{ pb}^{-1}$.

- Standard Model Higgs:
 $m_H > 107 \text{ GeV}$ at 95% C.L
- SUSY
- $M_{\tilde{\chi}_1^\pm} > 102.7 \text{ GeV}$ (for large s-neutrino masses.)
- No significant excess to S.M. predictions for Scalar Leptons in Standard SUSY Scalar Top and Bottom Quarks
- Search for exotica and new physics.
- No sign of existence of Heavy Neutral and Stable Charged Leptons.
 $m_L^\pm > 102.4 \text{ GeV}$ at 95% C.L
- No sign of existence of Excited Leptons
mass $\ell^* \geq 100 \text{ GeV}$ at 95% C.L

Any sign of new physics in 2000 L3 Data?

NOT YET

Need more Luminosity at highest possible energy