

# Search for Excited Fermions at HERA


Ali Sabetfakhri (ZEUS)  
University of Toronto



Representing the **H1** and **ZEUS** Collaborations

- Introduction*
- Search for Excited Electrons*
- Search for Excited Neutrinos*
- Search for Excited Quarks*
- Summary*

## Compositeness $\Rightarrow$ Excited Fermion



## Phenomenological Model

$$F^* \rightarrow F + V \quad \hookrightarrow q \bar{q}', l \bar{l}'$$

$$(V = \gamma, Z, W)$$

$$\mathcal{L}_{F^*F} = \frac{1}{\Lambda} \bar{F}_R^* \left[ f \cdot SU(2)_W + f' \cdot U(1)_Y + f_s \cdot SU(3)_C \right] F_L$$


Compositeness Scale :  $\Lambda \sim \mathcal{O}(\text{TeV})$

Free Parameters :  $f, f', f_s \sim \mathcal{O}(1)$

$$\sigma(ep \rightarrow F^* X) \sim \left(\frac{f}{\Lambda}\right)^2, \quad \left(\frac{f'}{\Lambda}\right)^2, \quad \left(\frac{f_s}{\Lambda}\right)^2$$

$e^+p$ data	$e^-p$ data
$\sqrt{s} = 300 \text{ GeV}$	$\sqrt{s} = 318 \text{ GeV}$
$\mathcal{L} = 38 \text{ pb}^{-1}$	$\mathcal{L} = 17 \text{ pb}^{-1}$
$e^*, \nu^*, q^*$	$\nu^*$

## Event Displays






**H1**

$\nu^* \rightarrow \nu\gamma$   
candidate

**ZEUS**

$q^* \rightarrow qW \rightarrow qe\nu$   
candidate





Channel	Topology	Studied by
<b>Excited Electron</b>		
$e^* \rightarrow e\gamma$	$e + \gamma + \text{high } E_t$	ZEUS, H1
$e^* \rightarrow eZ \rightarrow eq\bar{q}$	$e + 2 \text{ jets}$	ZEUS, H1
$e^* \rightarrow eZ \rightarrow ee^+e^-$	$3 e + \text{high } E_t$	H1
$e^* \rightarrow eZ \rightarrow e\nu\bar{\nu}$	$e + \text{high } \cancel{P}_t$	H1
$e^* \rightarrow eZ \rightarrow e\mu\bar{\mu}$	$e + 2 \mu$	H1
$e^* \rightarrow \nu W \rightarrow \nu q'\bar{q}$	$\text{high } \cancel{P}_t + 2 \text{ jets}$	ZEUS, H1
<b>Excited Neutrino</b>		
$\nu^* \rightarrow \nu\gamma$	$\gamma + \text{high } \cancel{P}_t$	ZEUS, H1
$\nu^* \rightarrow eW \rightarrow e\nu e$	$2 e + \text{high } \cancel{P}_t$	H1
$\nu^* \rightarrow eW \rightarrow eq'\bar{q}$	$\text{high } \cancel{P}_t + 2 \text{ jets}$	ZEUS, H1
$\nu^* \rightarrow eW \rightarrow e\mu\nu$	$e + \mu + \text{high } \cancel{P}_t$	H1
$\nu^* \rightarrow \nu Z \rightarrow \nu q\bar{q}$	$\text{high } \cancel{P}_t + 2 \text{ jets}$	ZEUS, H1
$\nu^* \rightarrow \nu Z \rightarrow \nu ee$	$2 e + \text{high } \cancel{P}_t$	H1
<b>Excited Quark</b>		
$q^* \rightarrow q\gamma$	$\gamma + \text{jet}$	ZEUS, H1
$q^* \rightarrow qZ \rightarrow qq'\bar{q}'$	$3 \text{ jets}$	H1
$q^* \rightarrow qZ \rightarrow qee$	$2 e + \text{jet}$	H1
$q^* \rightarrow qZ \rightarrow q\mu\bar{\mu}$	$2 \mu + \text{high } \cancel{P}_t$	H1
$q^* \rightarrow qW \rightarrow q\nu e$	$e + \text{jet} + \text{high } \cancel{P}_t$	ZEUS, H1
$q^* \rightarrow qW \rightarrow qq'\bar{q}$	$3 \text{ jets}$	H1
$q^* \rightarrow qW \rightarrow q\mu\nu$	$\mu + \text{jet} + \text{high } \cancel{P}_t$	H1

# Search For Excited Electron

## Signature of main channels


- ▷  $e^* \rightarrow e\gamma$ : 1 electron + 1 photon (high  $E_t$ )
- ▷  $e^* \rightarrow \nu W \rightarrow \nu q'\bar{q}$ : high  $\cancel{p}_t$  + 2 high  $E_t$  jets (with mass  $\sim M_W$ )
- ▷  $e^* \rightarrow eZ \rightarrow eq\bar{q}$ : 1 electron + 2 high  $E_t$  jets (with mass  $\sim M_Z$ )

## H1


Channel	Events	Background
$e^* \rightarrow e\gamma$ (Ela)	53	$52.5 \pm 1.5$
$e^* \rightarrow e\gamma$ (Inel)	68	$77.4 \pm 8.2$
$e^* \rightarrow eZ \rightarrow ee$	1	$0.9 \pm 0.4$
$e^* \rightarrow eZ \rightarrow \mu\bar{\mu}$	0	$0.35 \pm 0.05$
$e^* \rightarrow eZ \rightarrow \mu\bar{\nu}$	1	$2.7 \pm 0.4$
$e^* \rightarrow eZ \rightarrow q\bar{q}$	14	$12.3 \pm 3.4$
$e^* \rightarrow \nu W \rightarrow q'\bar{q}$	3	$3.3 \pm 0.6$

## ZEUS

Channel	Events	Background
$e^* \rightarrow e\gamma$	60	$68 \pm 2$
$e^* \rightarrow eZ \rightarrow q\bar{q}$	29	$26.7 \pm 0.9$
$e^* \rightarrow \nu W \rightarrow q'\bar{q}$	12	$14.3 \pm 0.9$



## ZEUS 96+97 Preliminary




# HERA Excited Electron Limits

$$(f = f')$$

No evidence for  $e^*$  production

95% C.L. Limits on  $f/\Lambda$



## H1

Assuming  $f/\Lambda = 1/M_{e^*}$   
 $50 < M_{e^*} < 223$  GeV  
 are excluded.

## ZEUS

Assuming  $f/\Lambda = 1/M_{e^*}$   
 $100 < M_{e^*} < 229$  GeV  
 are excluded.

# Search For Excited Neutrino

## Signature of main channels

- ▷  $\nu^* \rightarrow \nu\gamma$  : 1 photon + high  $\cancel{P}_t$
- ▷  $\nu^* \rightarrow eW \rightarrow eq'\bar{q}$  : 1 electron + 2 high  $E_t$  jets (with mass  $\sim M_W$ )
- ▷  $\nu^* \rightarrow \nu Z \rightarrow \nu q\bar{q}$  : high  $\cancel{P}_t$  + 2 high  $E_t$  jets (with mass  $\sim M_Z$ )


### H1

Channel	Events	Background
$\nu^* \rightarrow \nu\gamma$	2	$2.56 \pm 0.17 \pm 1.20$
$\nu^* \rightarrow eW \rightarrow eq'\bar{q}$	4	$7.74 \pm 1.38 \pm 1.63$


### ZEUS

Channel	Events	Background
$\nu^* \rightarrow \nu\gamma$	2	$1.8 \pm 0.2$
$\nu^* \rightarrow eW \rightarrow q'\bar{q}$	20	$15.0 \pm 1.5$
$\nu^* \rightarrow \nu Z \rightarrow q\bar{q}$	16	$12.3 \pm 0.6$

ZEUS 98+99 Preliminary  
( $\nu^* \rightarrow eW \rightarrow eq'\bar{q}$ )



ZEUS 98+99 Preliminary  
( $\nu^* \rightarrow \nu Z \rightarrow \nu q\bar{q}$ )




# HERA Excited Neutrino Limits

$$(f = -f')$$

No evidence for  $\nu^*$  production

95% C.L. Limits on  $f/\Lambda$



## H1

Assuming  $f/\Lambda = 1/M_{\nu^*}$   
 $50 < M_{\nu^*} < 150$  GeV  
 are excluded.

## ZEUS

Assuming  $f/\Lambda = 1/M_{\nu^*}$   
 $100 < M_{\nu^*} < 163$  GeV  
 are excluded.




# HERA Excited Neutrino Limits

( $f = f'$  :  $\nu^* \rightarrow \nu\gamma$  forbidden)

No evidence for  $\nu^*$  production

95% C.L. Limits on  $f/\Lambda$



## H1

Assuming  $f/\Lambda = 1/M_{\nu^*}$   
 $110 < M_{\nu^*} < 134$  GeV  
 are excluded.

## ZEUS

Assuming  $f/\Lambda = 1/M_{\nu^*}$   
 $100 < M_{\nu^*} < 133$  GeV  
 are excluded.

# Search For Excited Quark

## Signature of main channels



- ▷  $q^* \rightarrow q\gamma$  : 1 photon + 1 jet
- ▷  $q^* \rightarrow qW \rightarrow qq'\bar{q}''$  : 3 high  $E_t$  jets
- ▷  $q^* \rightarrow qW \rightarrow qe\nu$  : 1 electron + high  $p_t$  + 1 jet
- ▷  $q^* \rightarrow qZ \rightarrow qq'\bar{q}'$  : 3 high  $E_t$  jets

## H1

Channel	Events	Background
$q^* \rightarrow q\gamma$	35	$36 \pm 5$
$q^* \rightarrow qZ_{\rightarrow ee}$	0	$0.65 \pm 0.53$
$q^* \rightarrow qZ_{\rightarrow \mu\bar{\mu}}$	0	$0.35 \pm 0.05$
$q^* \rightarrow qZ_{\rightarrow q\bar{q}}$	32	$25.3 \pm 9.1$
$q^* \rightarrow qW_{\rightarrow e\nu}$	1	$1.10 \pm 0.35$
$q^* \rightarrow qW_{\rightarrow \mu\nu}$	3	$0.41 \pm 0.03$
$q^* \rightarrow qW_{\rightarrow q'\bar{q}}$	39	$45.3 \pm 17.3$

## ZEUS

Channel	Events	Background
$q^* \rightarrow q\gamma$	10	$15.3 \pm 1.5$
$q^* \rightarrow qW_{\rightarrow e\nu}$	3	$3.3 \pm 0.5$




# HERA Excited Quark Limits

$(f = f', f_s = 0)$

No evidence for  $q^*$  production

95% C.L. Limits on  $f/\Lambda$




## H1

Assuming  $f/\Lambda = 1/M_{q^*}$   
 $50 < M_{q^*} < 188$  GeV  
 are excluded.

## ZEUS

Assuming  $f/\Lambda = 1/M_{q^*}$   
 $100 < M_{q^*} < 203$  GeV  
 are excluded.

95% C.L. Limits on  $f$   
 (Assuming  $f = f'$  and  $\Lambda = M_{q^*}$ )



For  $f_s < 0.1$   
**HERA** is more competitive than **Tevatron**

## Summary



- Searches for excited fermions have been performed by the **H1/ZEUS** collaborations using  $e^+p$  and  $e^-p$  data.
- No evidence for excited fermion production is found.
- $e^-p$  data (despite smaller luminosity) has larger potential for  $\nu^*$  due to much higher  $\sigma$ .
- **HERA** extends limits beyond the reach of other colliders.
- $e^*$  and  $\nu^*$  explored well beyond LEP limits.
- $q^*$  complements Tevatron for **EW**  $q^*$  production and **HERA** is more competitive than Tevatron for  $f_s < 0.1$ .
- 2001-2006 upgraded **HERA** data ( $\sim 1 \text{ fb}^{-1}$ ) will bring much extended discovery potential (especially for  $\nu^*$ ).